

## Parathyroid Transplant Deceased Donor Criteria

Absolute Donor Exclusion Criteria for Parathyroid Transplantation		
#	Factor	Criteria
1	Malignancy	<ul style="list-style-type: none"> <li>• Donors with active cancer (donors receiving chemotherapy or radiation therapy or palliative cancer care within the past 5 years); except skin and prostate cancers (<i>see below</i>)</li> <li>• Primary brain tumour: grade III or IV glioblastoma with biopsy or debulking.</li> <li>• Donors with active or past history of melanoma</li> <li>• Thyroid cancer</li> </ul>
2	ABO	<ul style="list-style-type: none"> <li>• ABO incompatible</li> </ul>
3	Donor Type Age/Weight	<ul style="list-style-type: none"> <li>• DCD</li> <li>• NDD &lt; 16 and &gt; 60</li> </ul>
4	Infection / Communicable Diseases	<ul style="list-style-type: none"> <li>• Positive HIV, HbsAg, HTLV-I/II test results</li> <li>• Positive HCV NAT</li> <li>• West Nile Virus (WNV), or recent exposure to WNV</li> <li>• Rabies or within the last 6 months bitten by an animal and treated as if animal was rabid</li> <li>• Documented fungal septicaemia</li> <li>• Severe untreated sepsis; Sepsis with undermined etiology; MRSA sepsis</li> <li>• Untreated encephalitis of viral or of unknown etiology</li> <li>• Untreated meningitis of viral or of unknown etiology</li> <li>• Active disseminated tuberculosis</li> <li>• Ebola or known exposure to person with Ebola</li> <li>• Travelled to countries affected by Ebola within the past month</li> <li>• Diagnosis of COVID-19 with onset of symptoms or first positive test within 21 days</li> </ul>
5	Other	<ul style="list-style-type: none"> <li>• Polycystic Kidney Disease</li> <li>• On dialysis</li> <li>• Calcium level = hypercalcemia</li> <li>• Chronic kidney failure</li> <li>• Previous thyroidectomy or parathyroidectomy</li> <li>• Class 1 current positive virtual cross match</li> </ul>

Relative Donor Exclusion Criteria for Parathyroid Transplantation		
#	Factor	Criteria
1	Malignancy	<ul style="list-style-type: none"> <li>• Prostate Cancers may proceed under ExD if: <ul style="list-style-type: none"> <li>○ <i>No evidence of metastases</i></li> <li>○ <i>Gleason 6 PSA under 10 under active surveillance</i></li> <li>○ <i>If Gleason 7 or less and current PSA less than 2 treated by rad</i></li> <li>○ <i>If Gleason 7 and PSA undetectable after prostatectomy</i></li> </ul> </li> <li>• Avoid Gleason 8 or greater and all PSA over 20. PSAs between 10-20 is controversial and should be discussed with a urologist.</li> </ul>
2	Vaccine-Induced Thrombotic Thrombocytopenia (VITT)	<ul style="list-style-type: none"> <li>• Consult TSP-ID and CMO Transplant prior to proceeding if potential donor has a suspected diagnosis of vaccine-induced thrombotic thrombocytopenia (VITT)</li> </ul>